

THE RATIONAL COHOMOLOGY RING OF THE MODULI SPACE OF ABELIAN 3-FOLDS

RICHARD HAIN

1. INTRODUCTION

Suppose that R is a commutative ring with unit. Denote by $Sp_n(R)$ the group of automorphisms of R^{2n} that preserve the unimodular alternating form given by the matrix

$$\begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$$

In this note we compute the rational cohomology ring of $Sp_3(\mathbb{Z})$, or equivalently, of $\mathcal{A}_3$, the moduli space of principally polarized abelian 3-folds.

Denote the $\mathbb{Q}$ -Hodge structure of dimension 1 and weight $-2n$ by $\mathbb{Q}(n)$. (For those not interested in Hodge theory, just interpret this as one copy of $\mathbb{Q}$.) Denote by λ the first Chern class in $H^\bullet(\mathcal{A}_3; \mathbb{Q})$ of the Hodge bundle $\det \pi_* \Omega^1$ associated to the projection π of the universal abelian 3-fold to $\mathcal{A}_3$.

Theorem 1. *The cohomology groups of $\mathcal{A}_3$ are given by*

$$H^j(\mathcal{A}_3; \mathbb{Q}) \cong H^j(Sp_3(\mathbb{Z}); \mathbb{Q}) \cong \begin{cases} \mathbb{Q} & j = 0; \\ \mathbb{Q}(-1) & j = 2; \\ \mathbb{Q}(-2) & j = 4; \\ E & j = 6; \\ 0 & \text{otherwise,} \end{cases}$$

where E is a two-dimensional mixed Hodge structure which is an extension

$$0 \rightarrow \mathbb{Q}(-3) \rightarrow E \rightarrow \mathbb{Q}(-6) \rightarrow 0.$$

The ring structure is determined by the condition that $\lambda^3 \neq 0$.

I do not know whether the mixed Hodge structure (MHS) E on H^6 is split. Since $\mathcal{A}_3$ is a smooth stack over $\text{Spec } \mathbb{Z}$, I expect it to be a multiple (possibly trivial) of the class

$$\zeta(3) \in \mathbb{C}/i\pi^3\mathbb{Q} \cong \text{Ext}_{\mathcal{H}}^1(\mathbb{Q}, \mathbb{Q}(3))$$

given by the value of the Riemann zeta function at 3. Determining this class would be interesting.

As a corollary, we deduce the rational cohomology of $\overline{\mathcal{A}}_3$, the Satake compactification of $\mathcal{A}_3$.

Supported in part by grants from the National Science Foundation.

Theorem 2. *The rational cohomology ring of $\overline{\mathcal{A}}_3$ is given by*

$$H^j(\mathcal{A}_3; \mathbb{Q}) \cong \begin{cases} \mathbb{Q}(-n) & j = 2n, n \in \{0, 1, 2, 4, 5, 6\}; \\ B & j = 6; \\ 0 & \text{otherwise,} \end{cases}$$

where B is a 3-dimensional mixed Hodge structure which is an extension

$$0 \rightarrow \mathbb{Q}(0) \rightarrow B \rightarrow \mathbb{Q}(-3)^2 \rightarrow 0.$$

The ring structure is determined by the condition that the cohomology ring contain the graded ring $\mathbb{Q}[\lambda]/(\lambda^7)$, where λ has degree 2 and type $(1, 1)$.

Along the way, we compute the rational cohomology of $\overline{\mathcal{A}}_2$, the Satake compactification of $\mathcal{A}_2$ as well.

Proposition 3. *The rational cohomology ring of $\overline{\mathcal{A}}_2$ is given by*

$$H^\bullet(\overline{\mathcal{A}}_2; \mathbb{Q}) \cong \mathbb{Q}[\lambda]/(\lambda^4).$$

where λ is the first Chern class of the Hodge bundle

Denote the moduli space of smooth projective curves over the complex numbers by $\mathcal{M}_g$. Another consequence of the proof is the surjectivity of the homomorphism

$$H^\bullet(\mathcal{A}_3; \mathbb{Q}) \rightarrow H^\bullet(\mathcal{M}_3; \mathbb{Q})$$

induced by the period mapping $\mathcal{M}_3 \rightarrow \mathcal{A}_3$.

The computation of the rational cohomology of $\mathcal{A}_1$ is classical¹ and follows from the fact that the quotient of the upper half plane by $Sp_1(\mathbb{Z}) = SL_2(\mathbb{Z})$ is a copy of the affine line. The computation of the rational cohomology of $\mathcal{A}_2$ is (essentially) due to Igusa [11].² Brownstein and Lee [2, 3] have computed the integral cohomology of $Sp_2(\mathbb{Z})$.

Suppose that $g \geq 2$. Recall that the *mapping class group* Γ_g in genus g is the group of isotopy classes of orientation preserving diffeomorphisms of a closed, oriented surface S of genus g . Its rational cohomology is isomorphic to that of $\mathcal{M}_g$. The *Torelli group* T_g is defined to be the kernel of the natural homomorphism

$$(1) \quad \Gamma_g \rightarrow Sp(H_1(S; \mathbb{Z}))$$

where Sp denotes the symplectic group, and where $H_1(S; \mathbb{Z})$ is regarded as a symplectic module via its intersection form. Choosing a symplectic basis of $H_1(S; \mathbb{Z})$ gives an isomorphism $Sp_g(\mathbb{Z}) \cong Sp(H_1(S; \mathbb{Z}))$. One then obtains the well known extension

$$(2) \quad 1 \rightarrow T_g \rightarrow \Gamma_g \rightarrow Sp_g(\mathbb{Z}) \rightarrow 1.$$

The *extended Torelli group* $\widehat{T}_g$ is the preimage of the center $\{\pm I\}$ of $Sp(H_1(S; \mathbb{Z}))$ under (1). Equivalently, it is the group of isotopy classes of diffeomorphisms of S that act as $\pm I$ on $H_1(S)$. One has the extensions

$$1 \rightarrow T_g \rightarrow \widehat{T}_g \rightarrow \{\pm I\} \rightarrow 1$$

and

$$(3) \quad 1 \rightarrow \widehat{T}_g \rightarrow \Gamma_g \rightarrow PSp_g(\mathbb{Z}) \rightarrow 1$$

¹It is trivial except in degree 0.

²It is 1-dimensional in degrees 0 and 2, and trivial elsewhere.

where $PSp_g(\mathbb{Z})$ denotes the integral projective symplectic group $Sp_g(\mathbb{Z})/\{\pm I\}$.

Our approach to computing the cohomology of $Sp_3(\mathbb{Z})$ is to analyse the spectral sequence of the extension (3). This entails knowing the cohomology of Γ_3 (or equivalently, $\mathcal{M}_3$) and of $\widehat{T}_3$. Looijenga [12] computed the cohomology of $\mathcal{M}_3$ using the theory of Del-Pezzo surfaces:

$$H^j(\mathcal{M}_3; \mathbb{Q}) \cong \begin{cases} \mathbb{Q} & j = 0; \\ \mathbb{Q}(-1) & j = 1; \\ \mathbb{Q}(-6) & j = 6; \\ 0 & \text{otherwise.} \end{cases}$$

The cohomology of $\widehat{T}_3$ is computed in Section 3 using the stratified Morse Theory of Goresky and MacPherson [7]. We use their theory of non-proper Morse functions (see their Part II, Chapter 10) applied to the square of the distance to a point function restricted to the jacobian locus. This is an elaboration of a trick of Geoff Mess [13] which he used to show that the Torelli group in genus 2 is free of countable rank, and which was also used by Johnson and Millson (cf. [13]) to show that Torelli space in genus 3 does not have the homotopy type of a finite complex. Our use of Morse theory is analogous to Goresky and MacPherson's treatment [7, Part III] of complements of affine subspaces of euclidean spaces. [Look at Mess's paper.]

It seems to be a curious fact that in low genus ($g = 2, 3$ so far), it is easier to compute the rational cohomology groups of $\mathcal{M}_g$ than of $\mathcal{A}_g$. This is perhaps a reflection of the richness of curve theory — that it is a more powerful tool for understanding the geometry of $\mathcal{M}_g$ than the theory of abelian varieties is as a tool for understanding the geometry of $\mathcal{A}_g$. It will be interesting to know if this trend persists when $g \geq 4$, when $\mathcal{M}_g$ is no longer dense in $\mathcal{A}_g$. The cohomology of $\mathcal{M}_4$ is not yet known, nor does it seem tractable to compute the cohomology of the extended Torelli group in genus 4.

2. PRELIMINARIES

General references for this section are [8, Chapt. 3] and [9].

A *polarized abelian variety* is a compact complex torus A together with a cohomology class

$$\theta \in H^{1,1}(A) \cap H^2(A; \mathbb{Z})$$

whose translation invariant representative is positive. The corresponding complex line bundle is ample. The polarization θ can be regarded as a skew symmetric bilinear form on $H_1(A; \mathbb{Z})$. The polarization is *principal* if this form is unimodular.

Jacobians of curves are polarized by the intersection pairing on $H_1(C; \mathbb{Z}) \cong H_1(\text{Jac } C; \mathbb{Z})$. This form is unimodular, and so jacobians are canonically principally polarized abelian varieties.

A *framed* principally polarized abelian variety is a principally polarized abelian variety A together with a symplectic basis of $H_1(A; \mathbb{Z})$ with respect to the polarization θ .

Suppose that $g \geq 1$. The maximal compact subgroup of $Sp_g(\mathbb{R})$ is $U(g)$. The symmetric space $Sp_g(\mathbb{R})/U(g)$ is isomorphic to the rank g Siegel upper half space

$$\mathfrak{h}_g = \left\{ \begin{array}{l} \text{symmetric } g \times g \text{ complex matrices} \\ \text{with positive definite imaginary part} \end{array} \right\}$$

(See, [10], for example.) It has dimension $g(g+1)/2$. Taking a framed principally polarized abelian variety $(A; a_1, \dots, a_g, b_1, \dots, b_g)$ to the corresponding period matrix gives a bijection

$$(4) \quad \mathfrak{h}_g \cong \left\{ \begin{array}{l} \text{isomorphism classes of framed princi-} \\ \text{pally polarized abelian varieties} \end{array} \right\}$$

We will regard $\mathfrak{h}_g$ as the (fine) moduli space of framed principally polarized abelian varieties of dimension g .

Changing symplectic bases gives a natural left action of $Sp_g(\mathbb{Z})$ on the moduli space of framed principally polarized abelian varieties. There is clearly a natural left $Sp_g(\mathbb{Z})$ action on $\mathfrak{h}_g = Sp_g(\mathbb{Z})/U(g)$. The bijection (4) is equivariant with respect to these actions.

The moduli space $\mathcal{A}_g$ of principally polarized abelian varieties of dimension g is the quotient $Sp_g(\mathbb{Z}) \backslash \mathfrak{h}_g$. Since $\mathfrak{h}_g$ is contractible and since $Sp_g(\mathbb{Z})$ acts discontinuously and virtually freely on $\mathfrak{h}_g$, it follows that there is a natural isomorphism

$$H^\bullet(Sp_g(\mathbb{Z}); \mathbb{Q}) \cong H^\bullet(\mathcal{A}_g; \mathbb{Q}).$$

Taking a curve C to its jacobian $\text{Jac } C$ defines a morphism $\mathcal{M}_g \rightarrow \mathcal{A}_g$ which is called the *period mapping*.

Now suppose that $g \geq 2$. Denote Teichmüller space in genus g by $\mathcal{X}_g$. The mapping class group Γ_g acts properly discontinuously and virtually freely on $\mathcal{X}_g$ with quotient $\mathcal{M}_g$. It follows that there is a natural isomorphism

$$H^\bullet(\Gamma_g; \mathbb{Q}) \cong H^\bullet(\mathcal{M}_g; \mathbb{Q}).$$

We shall need several moduli spaces that sit between $\mathcal{X}_g$ and $\mathcal{M}_g$. Denote the quotient of $\mathcal{X}_g$ by T_g by $\mathcal{T}_g$. This space is known as *Torelli space*. Since $\mathcal{X}_g$ is contractible and T_g is torsion free, T_g acts freely on $\mathcal{X}_g$ and Torelli space is an Eilenberg-MacLane space with fundamental group T_g . Consequently,

$$H^\bullet(T_g; \mathbb{Z}) \cong H^\bullet(\mathcal{T}_g; \mathbb{Z}).$$

By a *framed Riemann surface* of genus g we shall mean a compact Riemann surface C together with a symplectic basis $a_1, \dots, a_g, b_1, \dots, b_g$ of $H_1(C; \mathbb{Z})$ with respect to the intersection form. Torelli space $\mathcal{T}_g$ is the moduli space of framed Riemann surfaces of genus g ; its points correspond to isomorphism classes of framed, genus g Riemann surfaces. The symplectic group $Sp_g(\mathbb{Z})$ acts on the framings in the natural way; the quotient $Sp_g(\mathbb{Z}) \backslash \mathcal{T}_g$ is $\mathcal{M}_g$.

Denote the locus in $\mathfrak{h}_g$ consisting of jacobians of smooth curves by $\mathcal{J}_g$. (Note that this is not closed in $\mathcal{A}_g$.) The period mapping $\mathcal{T}_g \rightarrow \mathcal{J}_g$ is surjective by definition. Since minus the identity is an automorphism of every polarized abelian variety A ,

$$(A; a_1, \dots, b_g) \cong (A; -a_1, \dots, -b_g).$$

But if C is a genus g curve, then

$$(C; a_1, \dots, b_g) \cong (C; -a_1, \dots, -b_g)$$

if and only if C is hyperelliptic. It follows that, when $g \geq 3$, the period mapping

$$\mathcal{T}_g \rightarrow \mathcal{J}_g$$

that takes a framed curve to its jacobian with the same framing is surjective and 2:1 except along the hyperelliptic locus, where it is 1:1. It follows that, when $g \geq 3$,

$$\mathcal{J}_g = \widehat{T}_g \backslash \mathcal{X}_g \text{ and } \mathcal{M}_g = PSp_g(\mathbb{Z}) \backslash \mathcal{J}_g.$$

The following diagram shows the coverings and their Galois group when $g \geq 3$:

$$(5) \quad \begin{array}{ccccccc} & & & \Gamma_g & & & \\ & & \hat{T}_g & & Sp_g(\mathbb{Z}) & & \\ & \nearrow & & \searrow & & \nearrow & \\ \mathcal{X}_g & \xrightarrow{T_g} & \mathcal{T}_g & \xrightarrow{\mathbb{Z}/2\mathbb{Z}} & \mathcal{J}_g & \xrightarrow{PSp_g(\mathbb{Z})} & \mathcal{M}_g \\ & & \searrow & & \downarrow & & \downarrow \\ & & & & \mathfrak{h}_g & \xrightarrow{PSp_g(\mathbb{Z})} & \mathcal{A}_g \end{array}$$

Lemma 4. *If $1 \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow E \rightarrow G \rightarrow 1$ is a group extension, then the projection $E \rightarrow G$ induces an isomorphism on homology and cohomology with 2-divisible coefficients. In particular*

$$H^\bullet(PSp_g(\mathbb{Z}); \mathbb{Z}[1/2]) \rightarrow H^\bullet(Sp_g(\mathbb{Z}); \mathbb{Z}[1/2])$$

is an isomorphism.

Proof. This follows from the fact that

$$H^j(\mathbb{Z}/2; \mathbb{Z}[1/2]) = 0 \quad j > 0$$

using the Hochschild-Serre spectral sequence of the group extension. $\square$

Proposition 5. *For all $g \geq 2$, there is a natural isomorphism*

$$H_\bullet(\mathcal{J}_g; \mathbb{Z}[1/2]) \cong H_\bullet(\hat{T}_g; \mathbb{Z}[1/2]) \cong H_\bullet(\mathcal{T}_g; \mathbb{Z}[1/2])^{\mathbb{Z}/2\mathbb{Z}}.$$

There are similar isomorphisms for cohomology.

Proof. Since T_g is torsion free, T_g acts fixed point freely on Teichmüller space, and $\mathcal{T}_g$ is a model of the classifying space of T_g . Recall that if X is a simplicial complex on which $\mathbb{Z}/2$ acts simplicially (but not necessarily fixed point freely), then the map

$$p_* : H_\bullet(X/(\mathbb{Z}/2); \mathbb{Z}[1/2]) \rightarrow H_\bullet(X; \mathbb{Z}[1/2])^{\mathbb{Z}/2}$$

induced by the projection p is an isomorphism, whose inverse is half the pullback map p^* . Applying this twice gives isomorphisms

$$H^\bullet(\mathcal{J}_g; \mathbb{Z}[1/2]) \cong H^\bullet(\mathcal{T}_g; \mathbb{Z}[1/2])^{\mathbb{Z}/2} \cong H^\bullet(T_g; \mathbb{Z}[1/2])^{\mathbb{Z}/2} \cong H^\bullet(\hat{T}_g; \mathbb{Z}[1/2]).$$

$\square$

A theta divisor of a principally polarized abelian variety A is a divisor Θ whose Poincaré dual is the polarization and which satisfies $i^*\Theta = \Theta$, where $i : x \mapsto -x$. Any two such divisors differ by translation by a point of order 2, and can be given as the zero locus of a theta function associated to a period matrix of A . A principally polarized abelian variety A is *reducible* if it is isomorphic (as a polarized variety) to the product of two proper abelian subvarieties. If $A = A_1 \times A_2$, then any theta divisor of A is reducible:

$$\Theta_A = (\Theta_{A_1} \times A_2) \cup (A_1 \times \Theta_{A_2}).$$

Denote the locus of reducible abelian varieties in $\mathcal{A}_g$ by $\mathcal{A}_g^{\text{red}}$ and in $\mathfrak{h}_g$ by $\mathfrak{h}_g^{\text{red}}$. Elements of $\mathfrak{h}_g^{\text{red}}$ are precisely those period matrices Ω that can be written as a direct sum of two smaller period matrices.

Proposition 6. *Denote the closure of $\mathcal{J}_g$ in $\mathfrak{h}_g$ by $\overline{\mathcal{J}}_g$. If $g \geq 2$, then $\mathcal{J}_g = \overline{\mathcal{J}}_g - (\overline{\mathcal{J}}_g \cap \mathfrak{h}_g^{\text{red}})$.*

Proof. The period mapping $\mathcal{M}_g \rightarrow \mathcal{A}_g$ extends to a morphism $\overline{\mathcal{M}}_g \rightarrow \overline{\mathcal{A}}_g$ from the Deligne-Mumford compactification of $\mathcal{M}_g$ to the Satake compactification of $\mathcal{A}_g$. The inverse image of the boundary $\overline{\mathcal{A}}_g - \mathcal{A}_g$ of $\overline{\mathcal{A}}_g$ is the boundary divisor Δ_0 of $\overline{\mathcal{M}}_g$, whose generic point is an irreducible stable curve of genus g with one node. Denote the moduli space of curves of compact type $\overline{\mathcal{M}}_g - \Delta_0$ by $\widetilde{\mathcal{M}}_g$. Since $\overline{\mathcal{M}}_g$ is complete, it follows that the period mapping $\widetilde{\mathcal{M}}_g \rightarrow \mathcal{A}_g$ is proper and therefore has closed image in $\mathcal{A}_g$. Since $\mathcal{M}_g$ is dense in $\overline{\mathcal{M}}_g$ and has image $Sp_g(\mathbb{Z}) \backslash \mathcal{J}_g$ under the period mapping, it follows that the image of $\widetilde{\mathcal{M}}_g$ in $\mathcal{A}_g$ is $Sp_g(\mathbb{Z}) \backslash \overline{\mathcal{J}}_g$.

Recall that the theta divisor $\Theta_C \subset \text{Jac } C$ of a smooth genus g curve C is (up to a translate by a point of order 2) the image of mapping

$$C^{g-1} \rightarrow \text{Pic}^{g-1} C \rightarrow \text{Jac } C$$

that takes $(x_1, \dots, x_{g-1})$ to $x_1 + \dots + x_{g-1} - \alpha$, where α is a square root of the canonical bundle of C . (See, for example, [8, p. 338].) It follows that Θ_C is irreducible. On the other hand, if C is a reducible, stable, curve of compact type, its jacobian is the product of the components of its irreducible components, and is therefore reducible. The result follows. $\square$

Corollary 7. *We have $\mathcal{J}_3 = \mathfrak{h}_3 - \mathfrak{h}_3^{\text{red}}$.*

Proof. Both $\mathcal{M}_3$ and $\mathcal{A}_3$ have dimension 6. Since the period mapping $\mathcal{M}_3 \rightarrow \mathcal{A}_3$ is generically of maximal rank, $\widetilde{\mathcal{M}}_3 \rightarrow \mathcal{A}_3$ is surjective. This implies that $\overline{\mathcal{J}}_3 = \mathfrak{h}_3$, from which the result follows. $\square$

3. THE HOMOLOGY OF $\mathcal{J}_3$ AND $\widehat{T}_3$

Denote the singular locus of an analytic variety Z by Z^{sing} . We shall compute the homology of $\mathcal{J}_3$ by applying stratified Morse theory to the stratification

$$(6) \quad \mathfrak{h}_3 \supseteq \mathfrak{h}_3^{\text{red}} \supset \mathfrak{h}_3^{\text{red, sing}}$$

of $\mathfrak{h}_3$. The top stratum is, by Corollary 7, $\mathcal{J}_3$. Note that there are natural inclusions

$$\mathfrak{h}_1 \times \mathfrak{h}_2 \hookrightarrow \mathfrak{h}_3^{\text{red}} \text{ and } \mathfrak{h}_1 \times \mathfrak{h}_1 \times \mathfrak{h}_1 \hookrightarrow \mathfrak{h}_3^{\text{red, sing}}$$

defined by

$$(\tau, \Omega) \mapsto \begin{pmatrix} \tau & 0 \\ 0 & \Omega \end{pmatrix} \text{ and } (\tau_1, \tau_2, \tau_3) \mapsto \begin{pmatrix} \tau_1 & 0 & 0 \\ 0 & \tau_2 & 0 \\ 0 & 0 & \tau_3 \end{pmatrix}$$

respectively. Note that the image of $\mathfrak{h}_1 \times \mathfrak{h}_2$ is stabilized in $Sp_3(\mathbb{R})$ by $SL_2(\mathbb{R}) \times Sp_2(\mathbb{R})$, and the image of $(\mathfrak{h}_1)^3$ by $\Sigma_3 \ltimes SL_2(\mathbb{R})^3$, where Σ_3 is identified with the subgroup

$$\{a_j \mapsto a_{\sigma(j)} \text{ and } b_j \mapsto b_{\sigma(j)} : \sigma \text{ is a permutation of } \{1, 2, 3\}\}$$

of $Sp_3(\mathbb{R})$. Here $a_1, \dots, b_6$ is the distinguished framing of the first homology of the corresponding abelian variety.

Proposition 8. *The stratification (6) satisfies Whitney's conditions (A) and (B) (cf. [7, p. 37]). Moreover*

$$(7) \quad \mathfrak{h}_3^{\text{red}} = \bigcup_{g \in Sp_3(\mathbb{Z})} g(\mathfrak{h}_1 \times \mathfrak{h}_2) = \bigcup_{g \in Sp_3(\mathbb{Z})/(SL_2(\mathbb{Z}) \times Sp_2(\mathbb{Z}))} g(\mathfrak{h}_1 \times \mathfrak{h}_2)$$

and

$$(8) \quad \mathfrak{h}_3^{\text{red}, \text{sing}} = \bigcup_{g \in Sp_3(\mathbb{Z})} g(\mathfrak{h}_1 \times \mathfrak{h}_1 \times \mathfrak{h}_1) = \coprod_{g \in Sp_3(\mathbb{Z})/(\Sigma_3 \ltimes (SL_2(\mathbb{Z}))^3)} g(\mathfrak{h}_1 \times \mathfrak{h}_1 \times \mathfrak{h}_1)$$

In particular, $\mathfrak{h}_3^{\text{red}}$ is a locally finite union of totally geodesic complex submanifolds of $\mathfrak{h}_3$ of complex codimension 2 and $\mathfrak{h}_3^{\text{sing}, \text{red}}$ is a countable disjoint union of totally geodesic complex submanifolds of $\mathfrak{h}_3$ of complex codimension 3.

Proof. Since every reducible abelian variety is the product (as polarized varieties) of an elliptic curve and an abelian surface, $\mathcal{A}_1 \times \mathcal{A}_2 \rightarrow \mathcal{A}_3^{\text{red}}$ is surjective. Lifting to $\mathfrak{h}_3$, this implies that the $Sp_3(\mathbb{Z})$ acts transitively on the components of $\mathfrak{h}_3^{\text{red}}$, and that $\mathfrak{h}_3^{\text{red}}$ is the $Sp_3(\mathbb{Z})$ -orbit of $\mathfrak{h}_1 \times \mathfrak{h}_2$ in $\mathfrak{h}_3$. Since $SL_2(\mathbb{R}) \times Sp_2(\mathbb{R})$ acts transitively on $\mathfrak{h}_1 \times \mathfrak{h}_2$, the stabilizer in $Sp_3(\mathbb{Z})$ of $\mathfrak{h}_1 \times \mathfrak{h}_2$ is $SL_2(\mathbb{Z}) \times Sp_2(\mathbb{Z})$. Assertion (8) follows.

The components of $\mathfrak{h}_3^{\text{red}}$ are smooth, so the $\mathfrak{h}_3^{\text{red}, \text{sing}}$ is the locus where two or more components of $\mathfrak{h}_3^{\text{red}}$ intersect. This is precisely the preimage of the locus in $\mathcal{A}_3$ of products of 3 elliptic curves. Since this locus is irreducible (it is the image of $(\mathcal{A}_1)^3 \rightarrow \mathcal{A}_3$), $\mathfrak{h}_3^{\text{red}, \text{sing}}$ is the $Sp_3(\mathbb{Z})$ -orbit of $(\mathfrak{h}_1)^3$. By the semi-simplicity of polarized abelian varieties, there is a unique way to decompose an element of $\mathfrak{h}_3^{\text{red}, \text{sing}}$ as a product of three elliptic curves. This, and the fact that $SL_2(\mathbb{R})^3$ acts transitively on $(\mathfrak{h}_1)^3$, imply the stabilizer in $Sp_3(\mathbb{Z})$ of $(\mathfrak{h}_1)^3$ is $\Sigma_3 \ltimes SL_2(\mathbb{Z})^3$ and that each component of $\mathfrak{h}_3^{\text{red}, \text{sing}}$ is smooth. This proves (7).

Whitney's condition (A) is automatic as each component of $\mathfrak{h}_3^{\text{red}, \text{sing}}$ is a homogeneous submanifold of the closure of each component of $\mathfrak{h}_3^{\text{red}}$. Condition (B) is well known to be a consequence of condition (A). $\square$

Integrating the Riemannian metric along geodesics gives an $Sp_g(\mathbb{R})$ -invariant distance function d on X . For a point $p \in X$, let $D_p : X \rightarrow \mathbb{R}$ be the square of the distance to p :

$$D_p(x) = d(x, p)^2.$$

The following result can be proved, either by appealing to [7, I.2.2.3] or by an elementary and direct argument.

Proposition 9. *There is an open dense subset U of X such that for all $p \in U$, $D_p : X \rightarrow \mathbb{R}$ is a Morse function in the sense of Goresky and MacPherson [7, p. 52], all of whose critical points are “nondepraved.”* $\square$

Since each stratum is a union of totally geodesic subspaces, and since the symmetric space metric is complete with non-positive curvature it follows that there is a unique critical point on each component of each stratum. Since the Morse data for each critical point is a product of the normal and tangential Morse data [7, p. 61], we only need compute the normal Morse data at each critical point. There are two types of these: those that lie on a translate of $\mathfrak{h}_1 \times \mathfrak{h}_2$ and those that lie on a translate of $(\mathfrak{h}_1)^3$. The Morse data at each depends only on its type.

Proposition 10. *If $x \in \mathfrak{h}_3^{\text{red}} - \mathfrak{h}_3^{\text{red,sing}}$ is a critical point of D_p , then the normal Morse data at x is homotopy equivalent to $(S^3, *)$. In particular, D_p is perfect at such critical points.*

Proof. Since the normal slice at a smooth point of $\mathfrak{h}_3^{\text{red}}$ is a complex 2-ball, the normal Morse data at x is $(S^3, *)$. Since $H_\bullet(S^3) \rightarrow H_\bullet(S^3, *)$ is surjective, D_p is perfect at x . $\square$

Lemma 11. *At each point of $\mathfrak{h}_3^{\text{red,sing}}$, there is a normal slice with coordinates (z_1, z_2, z_3) such that $\mathfrak{h}_3^{\text{red}}$ has three components with equations*

$$z_2 = z_3 = 0, \quad z_1 = z_3 = 0, \quad z_1 = z_2 = 0.$$

Proof. There are three obvious ways to deform the product $A = E \times E' \times E''$ of three elliptic curves, preserving the polarization, into $\mathfrak{h}_3^{\text{red}}$. Namely, one can deform one of the elliptic curves in $\mathfrak{h}_1$, and deform the product of the other two into $\mathfrak{h}_2 - \mathfrak{h}_2^{\text{red}}$. The semi-simplicity of abelian varieties implies that each component of $\mathfrak{h}_3^{\text{red,sing}}$ is smooth and there are no other ways to deform A into $\mathfrak{h}_3^{\text{red}}$. It follows that 3 components of $\mathfrak{h}_3^{\text{red}}$ intersect at each point of $\mathfrak{h}_3^{\text{red,sing}}$.




FIGURE 1. Local structure of $\mathfrak{h}_3^{\text{red}}$ near $\mathfrak{h}_3^{\text{red,sing}}$

Since each component of $\mathfrak{h}_3^{\text{red,sing}}$ is locally homogeneous, and since all are conjugate under the action of $Sp_3(\mathbb{Z})$, to determine the local structure of $\mathfrak{h}_3^{\text{red}}$ near $\mathfrak{h}_3^{\text{red,sing}}$, it suffices to write down local equations along the component $(\mathfrak{h}_1)^3$ of $\mathfrak{h}_3^{\text{red}}$. Here we can use the coordinates

$$\begin{pmatrix} \tau_1 & z_3 & z_2 \\ z_3 & \tau_2 & z_1 \\ z_2 & z_1 & \tau_3 \end{pmatrix}$$

where each $\tau_j \in \mathfrak{h}_1$, and (z_1, z_2, z_3) lies in a neighbourhood of the origin in $\mathbb{C}^3$ small enough to guarantee that this matrix has positive definite imaginary part. In these coordinates, the three components of $\mathfrak{h}_3^{\text{red}}$ have equations:

$$z_2 = z_3 = 0, \quad z_1 = z_3 = 0, \quad z_1 = z_2 = 0.$$

□

View S^5 as the unit sphere in $\mathbb{C}^3$. The intersection of each coordinate axis with S^5 is a linearly imbedded S^1 . Together these give an imbedding

$$k : S^1 \amalg S^1 \amalg S^1 \hookrightarrow S^5$$

where $\amalg$ denotes disjoint union. The boundary of a small tubular neighbourhood of each S^1 is a trivial S^3 bundle over S^1 . Taking one fiber of each and connecting them to an arbitrary point of $S^5 - (S^1 \cup S^1 \cup S^1)$ gives an imbedding

$$i : S^3 \vee S^3 \vee S^3 \hookrightarrow S^5.$$

Proposition 12. *If $x \in \mathfrak{h}_3^{\text{red,sing}}$ is a critical point of D_p , then the normal Morse data at x is homotopy equivalent to*

$$(S^5 - k(S^1 \amalg S^1 \amalg S^1), i((S^3 \vee S^3 \vee S^3))).$$

Proof. By the previous Lemma, the normal Morse data is the same as that for the distance squared to a point function for

$$\mathbb{C}^3 - \text{the union of the 3 coordinate axes.}$$

This is computed using [7, I.3.11.2)] as explained in [7, III.3.3)].

□

To compute the homology of $\mathcal{J}_3$, we need to compute the homology of each kind of Morse data. Denote the quotient of $\mathbb{Z}^3$ by the diagonal subgroup by V . It is isomorphic to $\mathbb{Z}^2$.

Proposition 13. *The homology of the Morse data at a critical point of $\mathfrak{h}_3^{\text{red,sing}}$ is*

$$H_j(S^5 - k(S^1 \amalg S^1 \amalg S^1), i(S^3 \vee S^3 \vee S^3); \mathbb{Z}) = \begin{cases} V & j = 4; \\ 0 & \text{otherwise.} \end{cases}$$

The homology is generated by the boundaries of tubular neighbourhoods of the three imbedded S^1 s, which are subject to the relation that their sum is zero. In particular, D_p is perfect at such critical points.

Proof. The computation is elementary. The Morse function is perfect at such critical points because the relative homology is generated by absolute cycles.

□

The natural action of the symmetric group Σ_3 on $\mathbb{Z}^3$ (by permuting the coordinates) preserves the diagonal subgroup and therefore descends to an action on V . We view V as a $\Sigma_3 \ltimes SL_2(\mathbb{Z})^3$ -module via the projection $\Sigma_3 \ltimes SL_2(\mathbb{Z})^3 \rightarrow \Sigma_3$.

Recall that if R is a commutative ring, K a subgroup of G , and M a RK module, then the G -module induced from M is defined by

$$\text{Ind}_K^G M := RG \otimes_{RK} M.$$

Theorem 14. *We have,*

$$H_j(\mathcal{J}_3; \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & j = 0; \\ \text{Ind}_{SL_2(\mathbb{Z}) \times Sp_2(\mathbb{Z})}^{Sp_3(\mathbb{Z})} \mathbb{Z} & j = 3; \\ \text{Ind}_{\Sigma_3 \ltimes SL_2(\mathbb{Z})^3}^{Sp_3(\mathbb{Z})} V & j = 4; \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Each critical point has trivial tangential Morse data, as the critical points are minima on each stratum. It follows from [7, I.3.7] that the Morse data at each critical point is homotopy equivalent to its normal Morse data. Since the Morse function is perfect, the homology of $\mathcal{J}_3$ is the sum of the relative homologies of the normal Morse data at each critical point. The action of $Sp_3(\mathbb{Z})$ follows from the description of the $Sp_3(\mathbb{Z})$ action on the strata in Proposition 8. $\square$

For a subgroup G of $Sp_g(\mathbb{Z})$ that contains $-I$, define PG to be the subgroup $G/(\pm I)$ of $PSp_g(\mathbb{Z})$.

Corollary 15. *For all $j \geq 0$,*

$$\begin{aligned} H_j(\widehat{T}_3; \mathbb{Z}[1/2]) &\cong H_k(T_3; \mathbb{Z}[1/2])^{\mathbb{Z}/2\mathbb{Z}} \cong \begin{cases} \mathbb{Z}[1/2] & j = 0; \\ \text{Ind}_{SL_2(\mathbb{Z}) \times Sp_2(\mathbb{Z})}^{Sp_3(\mathbb{Z})} \mathbb{Z}[1/2] & j = 3; \\ \text{Ind}_{\Sigma_3 \ltimes SL_2(\mathbb{Z})^3}^{Sp_3(\mathbb{Z})} V \otimes \mathbb{Z}[1/2] & j = 4; \\ 0 & \text{otherwise;} \end{cases} \\ &\cong \begin{cases} \mathbb{Z}[1/2] & j = 0; \\ \text{Ind}_{P(SL_2(\mathbb{Z}) \times Sp_2(\mathbb{Z}))}^{PSp_3(\mathbb{Z})} \mathbb{Z}[1/2] & j = 3; \\ \text{Ind}_{P(\Sigma_3 \ltimes SL_2(\mathbb{Z})^3)}^{PSp_3(\mathbb{Z})} V \otimes \mathbb{Z}[1/2] & j = 4; \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

4. THE SPECTRAL SEQUENCE

We shall compute the homology spectral sequence

$$(9) \quad H_i(PSp_3(\mathbb{Z}); H_j(\widehat{T}_3; \mathbb{Q})) \implies H_{i+j}(\Gamma_3; \mathbb{Q}).$$

Thanks to Shapiro's Lemma (see, for example, [1]) the E^2 term of the spectral sequence can be computed.

Lemma 16. *We have*

$$H_i(PSp_3(\mathbb{Z}); H_j(\widehat{T}_3; \mathbb{Q})) \cong \begin{cases} \mathbb{Q} & i = 0, 2 \text{ and } j = 3; \\ 0 & j > 0 \text{ and } j \neq 3. \end{cases}$$

Proof. Applying Shapiro's Lemma, Lemma 4, the Kunneth Theorem, and the fact that the rational homology of $SL_2(\mathbb{Z})$ is that of a point, we have:

$$\begin{aligned} H_i(PSp_3(\mathbb{Z}); \text{Ind}_{P(SL_2(\mathbb{Z}) \times Sp_2(\mathbb{Z}))}^{PSp_3(\mathbb{Z})} \mathbb{Q}) &\cong H_i(P(SL_2(\mathbb{Z}) \times Sp_2(\mathbb{Z})); \mathbb{Q}) \\ &\cong H_i(SL_2(\mathbb{Z}) \times Sp_2(\mathbb{Z}); \mathbb{Q}) \\ &\cong H_i(Sp_2(\mathbb{Z}); \mathbb{Q}). \end{aligned}$$

This is $\mathbb{Q}$ in when $i = 0, 2$ and 0 otherwise by Igusa's computation.

Let $V_{\mathbb{Q}} = V \otimes_{\mathbb{Z}} \mathbb{Q}$. This is the unique 2-dimensional irreducible representation of Σ_3 . Since $V_{\mathbb{Q}}$ is divisible and has no coinvariants, $H_{\bullet}(\Sigma_3; V_{\mathbb{Q}})$ vanishes in all degrees. Arguing as above, we have

$$\begin{aligned} H_i(PSp_3(\mathbb{Z}); \operatorname{Ind}_{P(\Sigma_3 \ltimes SL_2(\mathbb{Z})^3)}^{PSp_3(\mathbb{Z})} V_{\mathbb{Q}}) &\cong H_i(P(\Sigma_3 \ltimes SL_2(\mathbb{Z})^3); V_{\mathbb{Q}}) \\ &\cong H_i(\Sigma_3 \ltimes SL_2(\mathbb{Z})^3; V_{\mathbb{Q}}) \\ &\cong H_i(\Sigma_3; V_{\mathbb{Q}}) \\ &\cong 0. \end{aligned}$$

□

Proposition 17. *For $2 \leq r \leq 4$, the E^r -term of the spectral sequence (9) is*

4	0	0	0	0	0	0	0	0
3	$\mathbb{Q}$	0	$\mathbb{Q}$	0	0	0	0	0
2	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0
0	$\mathbb{Q}$	0	$\mathbb{Q}$	0	$\mathbb{Q}$	0	$\mathbb{Q}^2$	0
deg	0	1	2	3	4	5	6	7

(All terms not shown are zero.) In addition, the differentials

$$d^4 : E_{4,0}^4 \rightarrow E_{0,3}^4 \text{ and } d^4 : E_{6,0}^4 \rightarrow E_{2,3}^4$$

are both surjective, and $E^5 = E^\infty$.

Proof. The computation of $E_{s,t}^2$ for $t > 0$ follows from the previous lemma. It implies that $d^r = 0$ when $2 \leq r < 4$ and $E^5 = E^\infty$. By Looijenga's computation of the rational homology of Γ_3 we know that $H_3(\Gamma_3; \mathbb{Q})$ and $H_5(\Gamma_3; \mathbb{Q})$ both vanish. This implies that the differentials

$$d^4 : E_{4,0}^4 \rightarrow E_{0,3}^4 \text{ and } d^4 : E_{6,0}^4 \rightarrow E_{2,3}^4$$

must be surjective. Since $H_4(\Gamma_3; \mathbb{Q}) = 0$, and since $H_6(\Gamma_3; \mathbb{Q})$ is one dimensional, the first of these is an isomorphism and the second has one dimensional kernel. The result follows. □

Proof of Theorem 1. The computation of the rational homology (and therefore the rational cohomology) of $\mathcal{A}_3$ follows from Proposition 17.

Denote the first Chern class in $H^2(\mathcal{A}_3; \mathbb{Q})$ of the Hodge bundle by λ . It is the class of an ample line bundle. A standard argument, that uses the fact that the Satake compactification of A_3 has boundary of codimension 3, shows that there is a complete surface in A_3 . This implies that $\lambda^2 \neq 0$ in $H^4(\mathcal{A}_3; \mathbb{Q})$.

The proof that λ^3 does not vanish in $H^6(\mathcal{A}_3; \mathbb{Q})$ is more subtle, and is due to van der Geer [6]. We will give a topological proof of this fact in the next section. The key point in van der Geer's argument is that there is a complete subvariety of the characteristic p version $\mathcal{A}_{3/\mathbb{F}_p}$ of $\mathcal{A}_3$. This implies that λ^3 is not zero in $H_{\text{ét}}^6(\mathcal{A}_{3/\mathbb{F}_p}; \mathbb{Q}_\ell)$, where p is a prime where $\mathcal{A}_3$ has good reduction and ℓ is a prime distinct from p . Standard comparison theorems imply that this last group is isomorphic to $H^6(\mathcal{A}_3; \mathbb{Q}_\ell)$, which gives the desired non-vanishing of λ^3 .

The statement about weights follows as λ is of type $(1, 1)$. Since $\lambda^3 \neq 0$, this implies that $H^4(\mathcal{A}_3; \mathbb{Q})$ is generated by λ^2 and has type $(2, 2)$, and that $H^6(\mathcal{A}_3; \mathbb{Q})$

contains a copy of $\mathbb{Q}(-3)$ spanned by λ^3 . On the other hand, the spectral sequence in Proposition 17 implies that the restriction mapping

$$H^6(\mathcal{A}_3; \mathbb{Q}) \rightarrow H^6(\mathcal{M}_3; \mathbb{Q}) \cong \mathbb{Q}(-6)$$

is surjective, which completes the proof. $\square$

5. CYCLES

In this section, we give a topological proof that λ^3 is non-zero in $H^6(\mathcal{A}_3; \mathbb{Q})$. The approach is to construct a topological 6-cycle in $\mathcal{A}_3$ and then show that the value of λ^3 on it is non-zero.

Fix an imbedding of the Satake compactification $\overline{\mathcal{A}}_3$ of $\mathcal{A}_3$ in some projective space. Take a generic codimension 3 linear section of $\overline{\mathcal{A}}_3$ that avoids the boundary $\overline{\mathcal{A}}_3^{\text{red}} - \mathcal{A}_3^{\text{red}}$, is transverse to $\mathcal{A}_3^{\text{red}}$, and intersects $\mathcal{A}_3^{\text{red}, \text{sing}}$ transversally. This section is a complete curve X in $\mathcal{A}_3^{\text{red}}$, smooth (in the orbifold sense) away from its intersection with $\mathcal{A}_3^{\text{red}, \text{sing}}$. At each point x where it intersects $\mathcal{A}_3^{\text{red}, \text{sing}}$, it has three branches. Set $X' = X - (X \cap \mathcal{A}_3^{\text{red}, \text{sing}})$.





FIGURE 2.

The next step is to construct a 5-cycle in $\mathcal{M}_3$ which is an S^3 bundle over X away from the triple points, and where the three branches of this bundle at each triple point x are plumbed together using the normal Morse data at x .

Denote the moduli space of principally polarized abelian 3-folds with a level ℓ structure by $\mathcal{A}_3[\ell]$. This is the quotient of $\mathfrak{h}_3$ by the level ℓ subgroup $Sp_3(\mathbb{Z})[\ell]$ of $Sp_3(\mathbb{Z})$, and is smooth when $\ell \geq 3$. Fix an $\ell \geq 3$. The symmetric space metric on $\mathfrak{h}_3$ descends to $\mathcal{A}_3[\ell]$. With the help of the metric, the normal bundle of any stratum can be viewed as a subbundle of the tangent bundle of $\mathcal{A}_3[\ell]$. Using the exponential mapping, we can identify a neighbourhood of the zero section of the normal bundle as being imbedded in $\mathcal{A}_3[\ell]$. Let Y and Y' be the inverse images of X and X' in $\mathcal{A}_3[\ell]$.

Choose a positive real number ϵ such that the exponential mapping is an imbedding on the $\epsilon/4$ -ball B of the normal bundle of $\mathcal{A}_3^{\text{red}}[\ell]$ restricted to Y' , and also on the ϵ -ball B' of the normal bundle of $\mathcal{A}_3^{\text{red}, \text{sing}}[\ell]$ at each point of $Y - Y'$. Set $\tilde{D} = B \cup B'$ and $\tilde{W} = \partial\tilde{D}$. Denote their pushforwards to $\mathcal{A}_3$ by D and W . Then W is a 5-cycle in $\mathcal{M}_3$ which is generically an S^3 -bundle over X' . Note that D is a 6-chain in $\mathcal{A}_3$ with $\partial D = W$.

On the other hand, Looijenga's computation of the rational homology of $\mathcal{M}_3$, implies that W bounds a rational 6-chain E in $\mathcal{M}_3$. Set $Z = D - E$. This is a rational 6-cycle in $\mathcal{A}_3$. By construction, we have:

FIGURE 3. Picture of W near a triple point

Proposition 18. *The cycle Z intersects $\mathcal{A}_3^{\text{red}, \text{sing}}$ transversally (in the orbifold sense), and the intersection number of Z with $\mathcal{A}_3^{\text{red}, \text{sing}}$ is non-zero.* $\square$

Corollary 19. *The class of Z is non-trivial in $H_6(\mathcal{A}_3; \mathbb{Q})$ and the class of $\mathcal{A}_3^{\text{red}, \text{sing}}$ is non-trivial in $H^6(\mathcal{A}_3; \mathbb{Q})$.* $\square$

The proof of the non-triviality of λ^3 is completed by the following result.

Proposition 20. *The class of $\mathcal{A}_3^{\text{red}, \text{sing}}$ in $H^6(\mathcal{A}_3; \mathbb{Q})$ is a non-zero multiple of λ^3 .*

Proof. Denote the closure in $\overline{\mathcal{A}}_3$ of a subvariety X of $\mathcal{A}_3$ by $\overline{X}$. Set $\partial X = \overline{X} - X$. Since $\mathcal{A}_3$ is a rational homology manifold, the sequence

$$H_8(\overline{\mathcal{A}}_3^{\text{red}}, \partial \mathcal{A}_3^{\text{red}}; \mathbb{Q}) \rightarrow H^4(\mathcal{A}_3; \mathbb{Q}) \rightarrow H^4(\mathcal{M}_3; \mathbb{Q})$$

is exact. Since $\mathcal{A}_3^{\text{red}}$ is irreducible of dimension 4, the left hand group is one-dimensional and spanned by the fundamental class of $\mathcal{A}_3^{\text{red}}$. Since the middle group is one-dimensional and the right hand group trivial, we see that the class of $\mathcal{A}_3^{\text{red}}$ spans $H^4(\mathcal{A}_3; \mathbb{Q})$. On the other hand, since λ is ample, and $\mathcal{A}_3$ contains a complete surface, λ^2 also spans $H^4(\mathcal{A}_3; \mathbb{Q})$. It follows that there is a non-zero rational number c such that

$$\lambda^2 = c[\mathcal{A}_3^{\text{red}}] \text{ in } H^4(\mathcal{A}_3; \mathbb{Q}).$$

Denote the determinant of the Hodge bundle by $\mathcal{L}$. The class λ^3 is represented by the divisor of a section of the restriction of $\mathcal{L}$ to $\mathcal{A}_3^{\text{red}}$. This can be computed by pulling back along the mapping $\mathcal{A}_1 \times \mathcal{A}_2 \rightarrow \mathcal{A}_3^{\text{red}}$. Since the Picard group of $\mathcal{A}_1$ is torsion, we see that the pullback of $\mathcal{L}$ is represented mod-torsion by $\mathcal{A}_1 \times D$, where D is a cycle representing λ on $\mathcal{A}_2$. But the cusp form χ_{10} of weight 10 on $\mathcal{A}_2$ is a section of $\mathcal{L}^{10}$, and has divisor supported on $\mathcal{A}_2^{\text{red}}$, we see that λ is represented by a non-zero rational multiple of the cycle $\mathcal{A}_1 \times \mathcal{A}_2^{\text{red}}$ on $\mathcal{A}_1 \times \mathcal{A}_2$, and by a non-zero multiple of $\mathcal{A}_3^{\text{red}, \text{sing}}$ in $H^4(\mathcal{A}_3^{\text{red}}; \mathbb{Q})$. This implies that λ^3 is represented by a non-zero rational multiple of $\mathcal{A}_3^{\text{red}, \text{sing}}$ in $H^4(\mathcal{A}_3; \mathbb{Q})$. $\square$

6. THE RATIONAL COHOMOLOGY OF $\overline{\mathcal{A}}_2$ AND $\overline{\mathcal{A}}_3$

Note that $\overline{\mathcal{A}}_0$ is just a point. Suppose now that $g > 0$. Then

$$\overline{\mathcal{A}}_g = \mathcal{A}_g \amalg \overline{\mathcal{A}}_{g-1}.$$

Choose a triangulation of $\overline{\mathcal{A}}_g$ such that $\overline{\mathcal{A}}_{g-1}$ is a subcomplex. Let U_g be a regular PL neighbourhood of $\overline{\mathcal{A}}_{g-1}$ in $\overline{\mathcal{A}}_g$. Set $U_g^* = U_g - \overline{\mathcal{A}}_{g-1}$. This has the homotopy type of ∂U_g , which is a rational homology manifold (as $\overline{\mathcal{A}}_g[3]$ is smooth) of dimension $2d_g - 1$, where $d_g = g(g+1)/2$ is the dimension of $\mathcal{A}_g$. The cohomology of U_g^* has a mixed Hodge structure and the cup product

$$H^{k-1}(U_g^*; \mathbb{Q}) \otimes H^{2d_g-k}(U_g^*; \mathbb{Q}) \rightarrow H^{2d_g-1}(U_g^*; \mathbb{Q}) \cong \mathbb{Q}(-d_g)$$

is a perfect pairing of mixed Hodge structures (MHSs) (see [5], for example).

Since $\mathcal{A}_g$ is a rational homology manifold, Lefschetz duality gives an isomorphism

$$H_c^k(\mathcal{A}_g, \mathbb{Q}) \cong \text{Hom}(H^{2d_g-k}(\mathcal{A}_g), \mathbb{Q}(-d_g)).$$

The standard long exact sequence

$$(10) \quad \cdots \rightarrow H^{k-1}(U_g^*) \rightarrow H_c^k(\mathcal{A}_g; \mathbb{Q}) \rightarrow H^k(\mathcal{A}_g; \mathbb{Q}) \rightarrow H^k(U_g^*; \mathbb{Q}) \rightarrow \cdots$$

is exact in the category of MHSs. These facts will allow us to compute the cohomology of U_2^* and U_3^* .

The second step in the computation will be to use the Mayer-Vietoris sequence associated to the covering $\overline{\mathcal{A}}_g = \mathcal{A}_g \cup U_g$

$$(11) \quad \cdots \rightarrow H^k(\overline{\mathcal{A}}_g) \rightarrow H^k(\mathcal{A}_g) \oplus H^k(\overline{\mathcal{A}}_{g-1}) \rightarrow H^k(U_g^*) \rightarrow H^{k+1}(\overline{\mathcal{A}}_g) \rightarrow \cdots$$

associated to the covering $\overline{\mathcal{A}}_g = \mathcal{A}_g \cup U_g$, which is exact in the category of MHS, to compute the cohomology of $\overline{\mathcal{A}}_2$ and $\overline{\mathcal{A}}_3$.

For determining the ring structure and also for seeing that some maps in these long exact sequences are non-trivial, it is useful to note that since λ is the class of an ample line bundle on $\overline{\mathcal{A}}_g$, the rational cohomology ring of $\overline{\mathcal{A}}_g$ contains the ring $\mathbb{Q}[\lambda]/(\lambda^{d_g+1})$.

6.1. Proof of Proposition 3. Since $H^\bullet(\mathcal{A}_2; \mathbb{Q})$ is $\mathbb{Q}$ in degree 0, $\mathbb{Q}(-1)$ in degree 2, and 0 otherwise, $H_c^\bullet(\mathcal{A}_2; \mathbb{Q})$ is $\mathbb{Q}(-2)$ in degree 4, $\mathbb{Q}(-3)$ in degree 6, and 0 otherwise. Using the sequence (10) and the fact that the cohomology of U_2^* satisfies Poincaré duality, we have

$$H^j(U_2^*; \mathbb{Q}) \cong \begin{cases} \mathbb{Q} & j = 0; \\ \mathbb{Q}(-1) & j = 2; \\ \mathbb{Q}(-2) & j = 3; \\ \mathbb{Q}(-3) & j = 5. \end{cases}$$

Putting this into the Mayer-Vietoris sequence (11), and using the fact that $\overline{\mathcal{A}}_1$ is $\mathbb{P}^1$, we obtain the result.

6.2. Proof of Proposition 2. By duality, $H_c^\bullet(\mathcal{A}_3; \mathbb{Q})$ is an extension of $\mathbb{Q}(-3)$ by $\mathbb{Q}(0)$ in degree 6, $\mathbb{Q}(-4)$ in degree 8, $\mathbb{Q}(-5)$ in degree 10, and $\mathbb{Q}(-6)$ in degree 12. Using the sequence (10) and the facts that $H^j(\mathcal{A}_3; \mathbb{Q})$ vanishes when $j \geq 7$ and $H_c^j(\mathcal{A}_3; \mathbb{Q})$ vanishes when $j \leq 5$, we have

$$H^j(U_3^*; \mathbb{Q}) \cong \begin{cases} H^j(\mathcal{A}_3; \mathbb{Q}) & j < 5; \\ H_c^{j+1}(\mathcal{A}_3; \mathbb{Q}) & j \geq 7. \end{cases}$$

In degrees 5 and 6 we have the exact sequence

$$0 \rightarrow H^5(U_3^*; \mathbb{Q}) \rightarrow H_c^6(\mathcal{A}_3; \mathbb{Q}) \xrightarrow{\alpha} H^6(\mathcal{A}_3; \mathbb{Q}) \rightarrow H^6(U_3^*; \mathbb{Q}) \rightarrow 0.$$

It follows from [4, 8.2.2] that the image of α is all of $W_6 H^6(\mathcal{A}_3; \mathbb{Q})$, so that α is non-zero. Since the sequence is exact in the category of MHSs, it follows that $H^5(U_3^*; \mathbb{Q})$ is $\mathbb{Q}(0)$ and $H^6(U_3^*; \mathbb{Q})$ is $\mathbb{Q}(-6)$.

Since $\overline{\mathcal{A}}_3$ is projective and λ is the class of a projective imbedding, the restriction of λ^j to U_3 is non-zero when $j = 1, 2, 3$. It follows from this and the computations above that λ and λ^2 restrict to non-trivial classes in the rational cohomology of U_3^* .

Putting all of this into the Mayer-Vietoris sequence (11), easily gives the computation of $H^j(\overline{\mathcal{A}}_3; \mathbb{Q})$ when $j \neq 6$. The computation of $H^6(\overline{\mathcal{A}}_3; \mathbb{Q})$ follows as the sequence

$$0 \rightarrow H^5(U_3^*; \mathbb{Q}) \rightarrow H^6(\overline{\mathcal{A}}_3; \mathbb{Q}) \rightarrow H^6(\mathcal{A}_3; \mathbb{Q}) \oplus H^6(\overline{\mathcal{A}}_2; \mathbb{Q}) \rightarrow H^6(U_3^*; \mathbb{Q}) \rightarrow 0$$

is exact in the category of MHSs.

REFERENCES

- [1] K. Brown: *Cohomology of groups*, Graduate Texts in Mathematics, 87, Springer-Verlag, New York, 1982.
- [2] A. Brownstein, R. Lee: *Cohomology of the symplectic group $\mathrm{Sp}_4(\mathbb{Z})$ I. The odd torsion case*, Trans. Amer. Math. Soc. 334 (1992), 575–596;
- [3] A. Brownstein, R. Lee: *Cohomology of the symplectic group $\mathrm{Sp}(4, \mathbb{Z})$. II. Computations at the prime 2*, Michigan Math. J. 41 (1994), 181–208.
- [4] P. Deligne: *Théorie de Hodge, III*, Inst. Hautes Études Sci. Publ. Math. No. 44 (1974), 5–77.
- [5] A. Durfee, R. Hain: *Mixed Hodge structures on the homotopy of links*, Math. Ann. 280 (1988), 69–83.
- [6] G. van der Geer: *The Chow ring of the moduli space of abelian threefolds*, J. Algebraic Geom. 7 (1998), 753–770.
- [7] M. Goresky, R. MacPherson: *Stratified Morse Theory*, Springer-Verlag, 1988.
- [8] P. Griffiths, J. Harris: *Principles of Algebraic Geometry*, Wiley-Interscience, 1978.
- [9] R. Hain, E. Looijenga: *Mapping class groups and moduli spaces of curves*, in Algebraic geometry—Santa Cruz 1995, Proc. Sympos. Pure Math., 62, Part 2 (1997), 97–142.
- [10] S. Helgason: *Differential geometry, Lie groups, and symmetric spaces*, Pure and Applied Mathematics, 80. Academic Press, 1978.
- [11] J. Igusa: *On Siegel modular forms of genus two*, Amer. J. Math. 84 (1962), 175–200.
- [12] E. Looijenga: *Cohomology of $\mathcal{M}_3$ and $\mathcal{M}_3^1$* , Mapping class groups and moduli spaces of Riemann surfaces (Göttingen, 1991/Seattle, WA, 1991), Contemp. Math., 150, 1993, 205–228,
- [13] G. Mess: *The Torelli groups for genus 2 and 3 surfaces*, Topology 31 (1992), 775–790.

DEPARTMENT OF MATHEMATICS, DUKE UNIVERSITY, DURHAM, NC 27708-0320
E-mail address: hain@math.duke.edu